

COSMOPOLITAN BULRUSH


Schoenoplectus maritimus (L.) Lye
Plant Symbol = SCMA8

Common Names: alkali bulrush

Scientific Names: *Bulboschoenus maritimus*, *Scirpus maritimus*

Description

General: Sedge Family (Cyperaceae). Alkali bulrush is a native perennial, heavily rhizomatous, obligate, wetland plant that may reach 1.5 m (60 in) in height and form dense stands. The stems are upright and angular with several leaves, up to 1 cm (0.4 in) wide, along the lower two thirds of the plant. The flowers are borne in sessile spikelets, densely clustered at the tip of the stem, and nestled in 3 or more leafy bracts. Spikelets are 1.2-2 cm (0.5 to 0.8 in) long. The seeds are brown lenticular achenes, 2.5 to 4 mm (0.1 to 0.16 in) long (Cronquist et al., 1977).



Distribution:

Cosmopolitan bulrush is found throughout North America with the exception of the Southeastern United States. For current distribution, consult the Plant Profile page for this species on the PLANTS Web site.

Habitat:

Cosmopolitan bulrush is found in areas with saturated soils including marshes, playas, ponds, streams and mud flats (Welsh et al., 2003). It is often found growing in association with other salt tolerant wetland species such as iodine bush (*Allenrolfea occidentalis*) inland saltgrass (*Distichlis spicata*) and seepweed (*Suaeda* spp.).

Adaptation

Cosmopolitan bulrush is found at low to mid elevations from 850 to 2,100 m (2,800 to 6,900 ft). in marshes, transient wet spots, pond margins, and backwater areas. It forms large dense stands in alkaline or saline sites. It can handle a pH of up to 9.0 and will grow on soils from fine clay to silt loam to sand. Cosmopolitan bulrush can survive periods of total inundation of up to 1 m (3 ft) deep. It tends to spread and reproduce when the water table is within 10 cm (4 in) of the surface. This species can occur in freshwater sites, but is usually a pioneering species that will be replaced over time with more permanent species. Cosmopolitan bulrush is fairly resistant to fire, which will increase its production and protein content.

Uses

Erosion Control, Restoration, & Constructed Wetlands: As a pioneering species, cosmopolitan bulrush will provide protection from wind and wave erosion especially for newly exposed soil. The rhizomes form a matrix for many beneficial bacteria making this plant an excellent choice for wastewater treatment constructed wetlands (Hoag and Sellers, 1995).

Wildlife & Livestock: Livestock and big game will rarely use this species for food. Palatability is low.

Waterfowl will utilize the seed and use the stems for nesting cover. Muskrats and beaver eat the rootstocks and young shoots, and will also use the shoots for building material.

Status

Threatened or Endangered: Cosmopolitan bulrush is a Species of Special Concern in Connecticut and Rhode Island, and Endangered in Illinois, New Jersey and New York (USDA NRCS, 2017).

Wetland Indicator: OBL

Please consult the PLANTS Web site (<http://plants.usda.gov/>) and your state's Department of Natural Resources for this plant's current status (e.g., threatened or endangered species, state noxious status, and wetland indicator values).

Planting Guidelines

Best establishment comes from planting plugs (either from the greenhouse or wild transplants). Plug spacing of 30 to 45 cm (12 to 18 in) will fill in within one growing season. Soil should be kept saturated. It can handle from 5-8 cm (2 to 3 in) of standing water during the establishment year. Fluctuating water levels during the establishment period is essential. Water levels can be managed to enhance spread and control weeds (Hoag, et al., 1992).

Wild plants for transplant can be collected and transplanted directly into the desired site. As long as no more than 4 dm² is removed from any 1-m² area (1 ft² in 1 yd²), the hole will fill in within one growing season. Care should be taken not to collect plants from weedy areas, as these weeds can be relocated to the transplant site, and the hole left at the collection site may fill with undesirable species.

Management

Water level in constructed wetlands should be fluctuated from saturated conditions up to a maximum depth of 30 cm (12 in) of standing water for establishing plants. The young plants can handle deeper water, but not for an extended period of time. This species can tolerate periods of drought and total inundation. Water levels can be managed to either enhance or reduce spread as well as to control terrestrial weeds.

Pests and Potential Problems

Generally, insects and disease are not a significant problem. If an insect or disease problem is encountered in the greenhouse, treat as you would for any other type plant.

Environmental Concerns

Cosmopolitan bulrush is native to North America. It can spread under favorable conditions but does not pose any environmental concern to native plant communities.

Seed and Plant Production

Cosmopolitan bulrush reproduces sexually by seed and asexually via rhizomes.

Seed Collection: Seed ripens in late August to October. Seeds are held tightly in the seed head, which means the collection time can be extended. Seed is typically collected by hand stripping the seed from the plant or clipping it using a pair of hand shears. Power seed harvesters are also effective. The bracts, which are found in the seed heads, are very irritating to the skin. Gloves and protective eye wear should be worn, especially when using a power seed harvester.

Seed Cleaning: Hammer mills can be used to break up the large debris and knock the seed loose from the stem. Cleaning can be accomplished using a seed cleaner with a No. 8 round top screen and a 1/8-inch bottom screen. Screens should be sized so desired seed will fall through and debris and weed seed are removed. Air velocity should be adjusted so chaff is blown away. Air flow and screen size may require adjustment to optimize cleaning process for given situation.

Greenhouse Plant Production: Propagation: Stratifying the seed in a mixture of water and sphagnum moss at 2°C (35° F) for 30 days may enhance the germination rate. Seed viability is quite high if stored properly for up to 20 years.

Seeds need light, moisture, and heat for germination. Place seeds on surface of soil and press in lightly to assure good soil contact. Do not cover seed. Soil should be kept moist. Greenhouse temperatures of (32 to 38°C (90 to 100° F) are required for germination and good growth. Germination should begin within about one week. Maintain moisture until plants are to be transplanted (Hoag and Sellers, 1994).

Cultivars, Improved, and Selected Materials (and area of origin)

There are no cultivars, improved, or selected materials of cosmopolitan bulrush. Common wildland collected seed is available from commercial sources. Cultivars should be selected based on the local climate, resistance to local pests, and intended use. Consult with your local land grant university, local extension or local USDA NRCS office for recommendations on adapted cultivars for use in your area.

Literature Cited

Cronquist, A., A.H. Holmgren, N.H. Holmgren, J.L. Reveal, & P.K. Holmgren 1977. Intermountain flora. Volume Six. The monocotyledons. Columbia University Press, New York, New York. 584p.

Hoag, J.C. & M. Sellers 1994. Riparian/wetland project information series no. 8: Use of greenhouse propagated wetland plants versus live transplants to vegetate constructed or created wetlands. USDA, NRCS, Idaho Plant Materials Center, Aberdeen, Idaho. 4p.

Hoag, J.C. & M. Sellers 1995. Riparian/wetland project information series no. 7: Constructed wetland system for water quality improvement of irrigation wastewater. USDA, NRCS, Idaho Plant Materials Center, Aberdeen, Idaho. 8p.

Hoag, J.C., G.L. Young, & J. Gibb 1992. Riparian/wetland project information series no. 1: Planting techniques for vegetation riparian areas from the Aberdeen Plant Materials Center. USDA, NRCS, Idaho Plant Materials Center, Aberdeen, Idaho. 8p.

USDA-NRCS. 2017. PLANTS Database. <https://www.plants.usda.gov/core/profile?symbol=BOMA7>. Accessed January 30, 2017.

Welsh, SL, Atwood ND, Goodrich, S., and LC Higgins. 2003. A Utah Flora. Third Edition, revised. Brigham Young University, Provo, UT.

Citation

Tilley, D., and L. St. John. 2012. Plant guide for cosmopolitan bulrush (*Schoenoplectus maritimus*). USDA-Natural Resources Conservation Service, Idaho Plant Materials Center. Aberdeen, ID. 83210.
Published: October 2012

Edited: 13sep2012djt; 30jan2017djt

For more information about this and other plants, please contact your local NRCS field office or Conservation District at <http://www.nrcs.usda.gov/> and visit the PLANTS Web site at <http://plants.usda.gov/> or the Plant Materials Program Web site: <http://plant-materials.nrcs.usda.gov>.

PLANTS is not responsible for the content or availability of other Web sites.

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the bases of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, or all or part of an individual's income is derived from any public assistance program, or protected genetic information in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases will apply to all programs and/or employment activities.)

If you wish to file an employment complaint, you must contact your agency's [EEO Counselor](#) (PDF) within 45 days of the date of the alleged discriminatory act, event, or in the case of a personnel action. Additional information can be found online at http://www.ascr.usda.gov/complaint_filing_file.html.

If you wish to file a Civil Rights program complaint of discrimination, complete the [USDA Program Discrimination Complaint Form](#) (PDF), found online at http://www.ascr.usda.gov/complaint_filing_cust.html, or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter to us by mail at U.S. Department of Agriculture, Director, Office of Adjudication, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, by fax (202) 690-7442 or email at program.intake@usda.gov. Individuals who are deaf, hard of hearing or have speech disabilities and you wish to file either an EEO or program complaint please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

Persons with disabilities who wish to file a program complaint, please see information above on how to contact us by mail directly or by email. If you require alternative means of communication for program information (e.g., Braille, large print, audiotape, etc.) please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

For any other information dealing with Supplemental Nutrition Assistance Program (SNAP) issues, persons should either contact the USDA SNAP Hotline Number at (800) 221-5689, which is also in Spanish or call the [State Information/Hotline Numbers](#). For any other information not pertaining to civil rights, please refer to the listing of the [USDA Agencies and Offices](#) for specific agency information.

Helping People Help the Land

USDA IS AN EQUAL OPPORTUNITY PROVIDER AND EMPLOYER